MAUSAM, 76, 3 (July 2025), 807-824

MAUSAM

DOI : https://doi.org/10.54302/mausam.v76i3.6635 Homepage: https://mausamjournal.imd.gov.in/index.php/MAUSAM



UDC No. 502.3 :504.5 : 621.43.064

17-year trend of tropospheric columnar NO₂ over India's northeast region observed by OMI: investigating probable anthropogenic and natural sources

ARUP BORGOHAIN^{1*}, ARBAN S. YOUROI^{1#}, ROHIT GAUTAM^{1,2!}, MANASI GOGOI^{2%}, RIBANDA

MARBANIANG⁵, NILAMONI BARMAN⁴, ARUNDHATI KUNDU^{1,3}, ABHAY SRIVASTAVA^{1@},

SHYAM S. KUNDU^{1\$} and S.P. AGGARWAL^{1&}

¹North Eastern Space Applications Centre, Dept. of Space, Govt. of India, Meghalaya-793103 (#arbanyouroi@gmail.com; @dr.abhaysrivastava@nesac.gov.in; \$ssk.nesac@gmail.com;

&spaggarwal2010@gmail.com)

²Department of Physics, Dibrugarh University, Dibrugarh, Assam-786004

(%manasi.gogoi93@gmail.com; !rohitgtm62@gmail.com)

³Department of Physics, Cotton University, Guwahati, Assam-781001 (to_arundhati@outlook.com)

⁴Central Ground Water Board, Eastern Region, Kolkata, West Bengal-700091 (neelspinor@gmail.com)

⁵Gauhati University, Guwahati (ribandamarbaniang@gmail.com)

(Received 1 May 2024, Accepted 25 February 2025)

*Corresponding author's email: arupborgohain@gmail.com

सार — यह अध्ययन भारत के पूर्वोत्तर क्षेत्र (NERI) में 17 वर्षों (2005-2022) में फैले क्षोभमंडलीय स्तंभाकार NO2 (TCN) सांद्रता का व्यापक विश्लेषण प्रस्तुत करता है। औरा उपग्रह पर लगे ओजोन मॉनिटरिंग इंस्डूमेंट (OMI) से सुदूर संवेदन डेटा का उपयोग क्षेत्र में नाइट्रोजन डाइऑक्साइड (NO2) सांद्रता के स्थानिक-कालिक पैटर्न का विश्लेषण करने में किया गया। NO2 एक प्रमुख वायुमंडलीय प्रदूषक है जो औद्योगिक उत्सर्जन, वाहन दहन, जैवभार दहन और प्राकृतिक प्रक्रियाओं जैसे तड़ित और मृदा उत्सर्जन जैसे विविध स्रोतों से निकलता है। NERI में NO2 प्रदूषण के विभिन्न स्तरों के लिए इसकी विशिष्ट स्थलाकृति और मौसम संबंधी कारकों के साथ-साथ शहरीकरण, जनसंख्या वृद्धि और ऊर्जा उपयोग के लिए जिम्मेदार ठहराया जा सकता है। मॉनसून पूर्व और शीत ऋतु के दौरान TCN सांद्रता बायोमास दहन और मानवजनित गतिविधियों जैसे कारकों के कारण चरम पर होती है। दीर्घकालिक आंकड़ों से पता चलता है कि समग्र TCN में वृद्धि हुई है, जो बढ़ते वाहनों, औद्योगिक विस्तार और जनसंख्या घनत्व के बढ़ते प्रभावों को दर्शाती है। मासिक परिवर्तन मॉनसून पूर्व ऋतु के महत्व को इंगित करते हैं, जिसमें तड़ित और परिवहनित NO2 के कारण NO2 का स्तर बढ़ जाता है। वर्मो की आग, बायोमास जलाना और दहन इंजन प्राकृतिक और मानवजनित NO2 के प्रमुख स्रोत आवृत्ति वितरण विश्लेषण के परिणाम एनईआरआई राज्यों में वायु गुणवत्ता की अलग-अलग स्थिति दर्शाते हैं जभा चरान्दर वह जीत है। यहा के परिणाम एनईआरआई राज्यों के वायु गुणवत्ता की अलग-अलग स्थिति दर्शाते

हैं, तथा लगातार उच्च टीसीएन स्तर का अनुभव करने वाले क्षेत्रों में लक्षित हस्तक्षेप की आवश्यकता पर बल देते हैं। इसके अलावा इस अध्ययन में कोविड-19 महामारी के प्रभाव का आकलन किया गया तथा मॉनसून पूर्व ऋतु में लॉकडाउन के दौरान नाइट्रोजन डाइऑक्साइड (NO2) की सांद्रता में उतार-चढ़ाव की देखी गई। यह शोध NERI में बढ़ते नाइट्रोजन डाइऑक्साइड प्रदूषण से निपटने के लिए निगरानी और शमन रणनीतियों की आवश्यकता पर जोर देता है, वायु गुणवत्ता और व्यापक पर्यावरणीय स्वास्थ्य मुद्दों को संबोधित करता है तथा स्वस्थ जीवन परिस्थितियों के लिए स्विचारित उपायों की आवश्यकता बताता है।

ABSTRACT. This study presents a comprehensive analysis of Tropospheric Columnar NO_2 (TCN) concentrations spanning 17 years (2005-2022) in the Northeastern Region of India (NERI). Remote sensing data from the Ozone Monitoring Instrument (OMI) aboard the Aura satellite was utilized in analyzing the spatiotemporal patterns of nitrogen dioxide (NO₂) concentrations within the region. NO₂ is a prominent atmospheric pollutant that emerges from diverse sources like industrial emissions, vehicle combustion, biomass burning, and natural processes such as lightning and soil emissions. The varying levels of NO₂ pollution in the NERI, with its distinctive topography and meteorological behaviors, may be attributed to urbanization, population growth, and energy utilization. TCN concentrations peak during pre-monsoon and winter months, driven primarily by factors like biomass burning and anthropogenic activities. Longterm data reveals an overall TCN increase, reflecting growing influences from rising vehicles, industrial expansion, and population density. Monthly variations indicate the significance of the pre-monsoon season, characterized by elevated NO₂ levels influenced by lightning and transported NO₂. Forest fires, biomass burning, and combustion engines contribute as major sources of both natural and anthropogenic NO₂. Frequency distribution analysis results exhibit varying air quality statuses across NERI states, emphasizing the need for targeted interventions in regions consistently experiencing high TCN levels. Furthermore, the study assesses the impact of the COVID-19 pandemic, identifying fluctuations in NO₂ concentrations during lockdowns in pre-monsoon seasons. This research emphasizes the requirement for strong monitoring and mitigation strategies to combat increasing NO₂ pollution in the NERI, addressing air quality and broader environmental health issues, necessitating well-informed measures for healthier living conditions.

Key words- Nitrogen dioxide, North Eastern Indian Region, Anthropogenic source, Lightning.

1. Introduction

Nitrogen dioxide (NO₂) significantly influences the Earth's atmospheric radiative balance, altering oxidation capacity and chemistry and impacting the lifetime of greenhouse gases. It primarily emanates from industrial processes, vehicle combustion, biomass fuel and crop residue burning, and soil emissions (Richter & Burrows, 2002; Cheng et al., 2012). Natural sources include lightning and soil emissions. NO₂ contributes to degraded air quality, adversely affecting human health (Molina and Molina, 2004). It is a key pollutant with regulated national ambient air quality standards worldwide. High tropospheric NO₂ concentrations pose a threat due to spatiotemporal variability influenced by local emissions, seasons, land use land cover (LULC) changes, and meteorological conditions. Satellite remote sensing is a viable approach. Studies in coal mining areas indicated positive correlations between thermal power plant output and NO_2 concentrations (Prasad *et al.*, 2012). Thunderstorm lightning has been considered a major source of nitrogen oxides (NOx, i.e., NO (nitric oxide) and NO₂) since von Liebig (1827) proposed it as a natural mechanism for the fixation of atmospheric nitrogen (Hutchinson, 1954). Lightning-induced nitrogen oxides (LNOx) have several important implications for atmospheric chemistry and climate (WMO, 1999; IPCC, 2001). Tropospheric (or ground-level) ozone also plays a role in the production of NO₂ through photochemical processes (Bradshaw et al., 1999) The global LNOx source is one of the largest natural sources of NOx in the atmosphere (Galloway et al., 2004) and certainly the largest source of NOx in the upper troposphere, in particular, in the tropics (WMO, 1999)

Noteworthy, NO₂ is also an important contributor to air pollution as a primary pollutant and as a precursor to ozone and fine particulate matter production. Human exposure to elevated NO₂ concentrations is associated with a range of adverse outcomes such as respiratory infections (Pannullo *et al.*, 2017; Tao *et al.*, 2014; Zeng *et al.*, 2020), increases in asthma incidence (Anenberg *et al.*, 2018; Achakulwisut *et al.*, 2019), lung cancer (Hamra *et al.*, 2015) and overall mortality (Brook *et al.*, 2007; Crouse *et al.*, 2015). NO₂ observations indicate air quality relationships with combustion sources of pollution such as transportation (Achakulwisut et al., 2019). Initial investigations found substantial decreases in the atmospheric NO₂ column from satellite observations (Goldberg et al., 2020; Biswal et al., 2021; Koukouli et al., 2021; Field et al., 2020; Bauwens et al., 2020; Liu et al., 2020; Prunet et al., 2020) and in ambient NO₂ concentrations from ground-based monitoring (Shi et al., 2021; Ropkins et al., 2021; Fu et al., 2020; Venter et al., 2020) during lockdowns enacted to reduce the spread of COVID-19. However, questions remain about the relationship of atmospheric columns with health- and policy-relevant ambient ground-level concentrations, and about the representativeness of sparse ground-based monitoring for broad assessment. Thus, there is a need to relate satellite observations of NO2 columns to groundlevel concentrations. It is also important to consider the effect of meteorology on recent NO₂ changes (Shi et al., 2021) and to quantify NO₂ changes due to COVID-19 interventions in the context of longer-term trends (Liu et al., 2021). Furthermore, air quality monitoring sites tend to be preferentially located in higher-income regions, raising questions about how NO2 changed in lowerincome regions where larger numbers of potentially susceptible people reside. Estimates of changes in groundlevel NO2 concentrations derived from satellite remote sensing would fill gaps between ground-based monitors, offer valuable information in regions with sparse and more clearly connect satellite monitoring, observations with ground-level ambient air quality.

North Eastern Region (NER) of India, Bangladesh, Bhutan, Tibetan Plateau, and Indo-Gangetic Plane (IGP) have a complex topography. These areas are affected by dry, cold westerly converge with hot moist southerly and help to develop thunderstorms associated with lightning (Kumar and Kamra 2012; Qie *et al.*, 2022). Lightning flashes are significantly correlated with convective rain, total column water vapor (TCWV), or surface relative humidity over both land and sea regions, according to previous studies (Price and Federmesser, 2006; Siingh *et al.*, 2011; Shi *et al.*, 2018). In this study, we utilized the tropospheric NO₂ and the lighting count over the North Eastern Region of India (NERI) to investigate the evolution and concentration trends across different states



Fig. 1. Topographical map of NERI adopted from topographic-map.com, the black lines represent the transportation of pollutants and the arrowhead indicates the direction

of the region. Specifically, we aim to quantify the increasing/decreasing trends of troposphere column NO2 and examine seasonal variations. Besides, the lightning count is also considered to connect with the NO₂ for its contribution. The NERI faces increasing NO₂ production due to population growth, urbanization, industrialization, agriculture demands, and energy consumption. Additionally, changes in LULC, such as deforestation and agricultural expansion, play a significant role in altering NO₂ levels by influencing local emissions and atmospheric dynamics. Assessing NO₂'s spatiotemporal distribution and identifying emission sources are crucial to formulating effective reduction strategies.

2. Data, Methodology and Study Area

2.1. Study domain and synoptic meteorology

The study area encompasses latitude 20° N - 30° N and longitude 88° E - 98° E, with a primary focus on the NERI. The NERI comprises of eight states: Assam (AS), Arunachal Pradesh (AR), Manipur (MN), Meghalaya (ML), Mizoram (MZ), Nagaland (NG), Tripura (TR) and Sikkim (SK) (Fig. 1). Notably, approximately 23.75% of India's forest area is found in the NERI (FSI, 2021). The region experiences a sub-tropical climate due to its distinctive topography. The Himalayan Mountains lie to the north, the Indo-Myanmar ranges of foothills and hills are in the east, and Mizoram and Tripura are to the south (Fig. 1). During the winter (Dec-Feb) and pre-monsoon (Mar-April) seasons, westerly winds significantly influence the atmospheric processes in the NER, as the region opens westward towards the IGP. The southwest monsoon serves as the primary source of rainfall in the area, and Cherrapunjee receives the highest rainfall globally, owing to orographic lifting by the Meghalaya plateau. The mean temperatures vary from 15 °C (January) to 28 °C (August) in the plain areas and from 9 °C to 21 °C in the hilly regions during the same periods. Relative humidity averages between 30-50% in the premonsoon season and 80-95% during the monsoon. To understand the wind patterns, we analyzed the averaged wind vector at 850 hPa from the European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysisversion 5 (ERA-5) over 15 years (2006-2020) from January to April, covering the area between 20° N - 30° N and 84° E-98° E. The analysis revealed dominant westerly winds during these months, with wind speeds ranging from 2 ms⁻¹ to 3 ms⁻¹ in January and February and 4 ms⁻¹ to 5 ms⁻¹ in March and April. Notably, during January and February, the westerly wind shifts to southwesterly over the Brahmaputra valley and further converts to southeasterly over the eastern part of the NER. Additionally, it converts to southwesterly/southeasterly over Nagaland, parts of Manipur, and other eastern borders.

2.2. Data Sources

Tropospheric NO₂ concentrations were obtained from the Ozone Monitoring Instrument (OMI) on the Aura satellite (Schoeberl *et al.*, 2004). Daily daytime global NO₂ datasets were generated by OMI aboard the polarorbiting, sun-synchronous Earth Observing System Aura satellite, with an equator crossing time of 13:45 LT. OMI measures the Earth's backscattered radiation within the spectral range of 264-504 nm (Streets *et al.*, 2013). To derive tropospheric column concentrations of NO₂, the



Fig. 2. Population, factories and vehicle growth over NERI

attenuation measurements provide slant column densities, converted to vertical column densities using appropriate air mass factors. Retrieval errors for NO2 in high-emission areas are up to 30% (Streets et al., 2013). The retrieval clear and cloudy process considers conditions, incorporating air mass factors derived from simulated NO₂ profiles (Bucselaet al., 2006). Tropospheric NO₂ has an uncertainty of 0.1×10^{15} mol.cm⁻² and is underestimated by 15-30% (Celarieret al., 2008). Land Use/Land Cover (LULC) data were obtained from ISRO's Bhuvan geoportal (https://bhuvan.nrsc.gov.in) collected using multi-temporal satellite imagery from the Resourcesat-1 LISS III sensor. Tropospheric ozone (O₃) concentrations were acquired from AURA OMI/MLS, computed by subtracting MLS-derived stratospheric ozone from OMI total column ozone (Ziemke et al., 2006). NO2 data was utilized which had been monitored as part of the National Air Quality Monitoring Programme (NAMP), which measures air pollutants including SO₂, NO₂ and PM₁₀ at 931 stations across 398 cities/towns in India. The monitoring is being carried out by Central Pollution Control Board (CPCB) in collaboration with other state and national agencies (https://cpcb.nic.in). Fire count datasets were utilized from a recent study (Borgohain

et al., 2023) of the impact of biomass burning over the region.

In this study, the Mann-Kendall test and Sens Slope are utilized for the trend analysis. The details about the technique are elaborated elsewhere (Theil, 1950; Sen 1968; Gilbert, 1987; Sirois, 1998; Wang and Swail, 2001; Yue *et al.*, 2002; Wang *et al.*, 2015).

2.3. Anthropogenic forcing in NERI

Based on data published by the Ministry of Statistics and Project Implementation, Government of India (www.mospi.gov.in) and projected population data from the Ministry of Health and Family Welfare, Government of India (GOI; https://nhm.gov.in), along with industrial the Survey data from Annual of Industries (http://microdata.gov.in) and vehicle from the data Transport Ministry of Road and Highways (https://morth.nic.in), Fig. 2 illustrates population growth, year-wise vehicle registrations, and factories over NERI from 2005 to 2019. Population data is available up to 2011 (As per GOI standard, there is a census every ten years, but since the 2021 census is still incomplete, projected



Fig. 3. Spatial distribution of mean TCN over the NERI (2005-2022)

data was only published). According to the combined census and projected data, Meghalaya experienced the largest population growth (~41%) in NERI, while Nagaland had the lowest (~10%). Overall population growth in NERI is around 31%, slightly below the national rate of about 32%. The number of vehicles and factories in NERI significantly increased, with data available up to 2019 and 2020, respectively. Vehicle numbers rose from 1.3 million to 6 million between 2005 and 2019, at 0.33 million per year. Factories increased from roughly 2200 to 6800 from 2005 to 2020, at 306 factories per year across NERI.

3. Results and discussion

3.1. Intra-annual variation over NERI

This study utilizes Tropospheric Columnar NO₂ (TCN) concentration data obtained from the Ozone Monitoring Instrument (OMI) spanning 17 years, from 2005 to 2022. Fig. 3 illustrates the temporal and spatial mean seasonal TCN over the eight states of NERI. The four seasons considered are pre-monsoon (March, April, May), monsoon (June, July, August, September), postmonsoon (October, November), and winter (December,

January, February). The annual mean TCN exhibits the highest concentrations in Assam $(1.57 \times 10^{15} \text{ mol cm}^{-2})$, Meghalaya $(1.60 \times 10^{15} \text{mol} \text{ cm}^{-2}),$ and Tripura $(1.62 \times 10^{15} \text{mol cm}^{-2})$, while Sikkim records the lowest (5.28×10¹⁴ mol cm⁻²). TCN levels are generally elevated during pre-monsoon and winter across all states, followed by post-monsoon and monsoon seasons. In the premonsoon period, Manipur shows the highest TCN (2.58×10¹⁵mol cm⁻²), followed by Mizoram and other states, while Arunachal Pradesh and Sikkim exhibit the lowest TCN (8.74×10¹⁴ mol cm⁻² and 5.27×10¹⁴ mol cm⁻ ²), respectively. This observed pattern of high TCN during the pre-monsoon months is attributed to significant biomass burning in these states (Borgohain et al., 2023).

The spatial and temporal distribution of TCN depends on meteorological factors, such as rainfall, wind direction, and Atmospheric Boundary Layer (ABL) height. TCN concentrations peak during the pre-monsoon months, followed by winter in all states, with significant contributions from biomass burning. The northeast part of India experiences elevated TCN levels during the premonsoon season due to extensive biomass burning emissions. Manipur (2.58×10¹⁵mol cm⁻²) and Mizoram $(2.37 \times 10^{15} \text{mol cm}^{-2})$ exhibit the highest TCN, while Sikkim and Arunachal Pradesh consistently record the lowest in all seasons. The pre-monsoon pattern of maximum NO₂ concentrations can be attributed to low humidity and mild temperatures, reducing the photolysis removal process of NO₂, thus stabilizing its concentration. Lower concentrations of OH (due to low humidity) also limit NO₂ photolysis and the formation of HNO₃, a principal sink for NO₂ (Jacob, 1999). In winter, high TCN is due to extensive biomass fuel usage for home heating and reduced availability of UV radiation to initiate photolysis reactions that break down NO₂ (Richter et al., 2005; Tariq et al., 2014; Uno et al., 2007; Nishanth and Kumar, 2011).

Additionally, a shallower ABL in winter results in lower vertical dispersion, reducing dilution and removal rates of NO₂, contributes to its enhancement (Barnes *et al.*, 2013). Elevated NO₂ concentrations during winter may also result from NO₂ transportation from the IGP to NERI due to westerly winds. Haridas *et al.* (2019) reported that the IGP shows the highest levels of NO₂ concentrations during winter.

Low TCN concentrations during monsoon result from wet removal processes. The presence of a moist clean air mass, increased actinic fluxes enhancing NO₂ photo dissociation, elevated OH radical levels aiding NO₂ removal via HNO₃, and potentially reduced traffic due to decreased social and educational activities during the hot summer contribute to the decrease in TCN (Ghude *et al.*, 2009; Tariq *et al.*, 2014; Ravindra *et al.*, 2003; Yoo *et al.*, 2014). During the rainy season, the impact of lightning on increasing NO₂ concentrations is not observed due to the opposing influence of rain washout (Yoo *et al.*, 2014).

3.2. Long-term TCN over NERI

Fig.4 illustrates the annual variations of TCN from 2005-2022 for four seasons as observed by the satellite over the eight northeastern states of India. Across all seasons, Assam consistently displayed the highest TCN value $(1.57 \times 10^{15} \text{mol cm}^{-2})$, along with Meghalaya, Tripura, and Nagaland. On the other hand, Sikkim and Arunachal Pradesh recorded the lowest TCN values $(5.28 \times 10^{14} \text{mol cm}^{-2})$. Notably, all states experienced an increase in TCN over the seventeen years (2005 to 2022) (Fig. 7). This long-term TCN increase is influenced by anthropogenic factors, such as rising vehicle numbers, industries, and population density.

Mann Kendall test and Sen Slope have been utilized for the TCN trend analysis and rate of change of TCN over the states. Arunachal Pradesh, Manipur, Nagaland, and Meghalaya show no trend in any season. Positive trends are detected in Assam during pre-monsoon and monsoon with Sen Slopes of 2.57×10¹³ and 1.65×10¹³, respectively. Pre-monsoon TCN increase is higher than during monsoon. Assam's valley region, developed cities, and national highways contribute to continuous anthropogenic emissions. Pre-monsoon TCN includes local and transported NO₂, while monsoon is dominated by locally emitted NO₂ due to the absence of transported NO₂. Meghalaya exhibits positive trends in winter and monsoon with Sen Slopes of 1.33×10^{13} and 1.66×10^{13} , respectively. Mizoram shows positive trends in monsoon and post-monsoon with Sen's Slopes of 1.23×10¹³and 9.96×10¹². Tripura displays positive trends in winter, monsoon and post-monsoon with Sen's Slopes of 1.99×10^{13} , 1.76×10^{13} and 3.38×10^{13} , respectively. The post-monsoon TCN increase is higher than in winter and monsoon. In winter and pre-monsoon, an aerosol river from North-west India reaches the NER corridor and bifurcates into two streams: one flows along the Brahmaputra valley (Assam), and the other flows along West Bengal and Bangladesh, finally reaching Tripura and Mizoram (south-east region). Kundu et al. (2018) reported that the Brahmaputra River valley in NERI is significantly affected by the prolonged dispersion of pollutants from the IGP during local winter and is burdened with a lot of natural and anthropogenic aerosol. Short-term trends for 2005-2008 showed increased NO₂ levels in the IGP, with some cities exhibiting positive NO₂-O₃ correlations (Streets et al., 2013). The IGP experienced an increasing NO₂ trend of 3×10^{13} mol/cm²/yr during 2005-2014 (Haridas et al., 2019). Ul - Haq et al. (2015) reported an



Fig. 4. State-wise seasonal TCN over NERI (2005-2022)



Fig. 5. Average monthly variation of NO₂ over different states of NER

average of $1.0 \pm 0.05 \times 15$ mol. cm⁻² with a 14% decadal increase in South Asia, linked to anthropogenic emissions from power generation, urbanization, vehicles, and industries.

High concentrations of NO_2 observed during winter may result from the transportation of NO_2 from the IGP to NERI due to westerly winds. Haridas *et al.* (2018) also reported that the IGP shows the highest NO_2 concentrations during winter. Road transport remains the dominant source of NO₂ emissions, with the number of registered vehicles in NERI increasing from 1.4 million in 2001 to approximately 6.1 million by 2019 (https://morth.nic.in). Another significant source of NO₂ emissions is industries, with the number of industries growing from 2200 to 6800 between 2005 and 2020. Population growth is a key factor contributing to the increase of TCN. About three in four rural households in India rely on traditional energy sources for cooking, heating, etc. Census 2011 shows that over 65 percent of



Fig. 6. Average seasonal variation of lightning flashes over different states of NERI

the population resides in rural India and consumes around 45 percent of the total domestic energy. While the use of modern energy sources is increasing, rural India continues to be predominantly dependent on traditional sources (Ranjan and Singh, 2017). Despite this, NERI generates significant agricultural waste, leading to NO₂ generation. In India, the annual crop residue generated for 2008-2009 was about 620 Mt/year, with 15% being burnt on farms, emitting substantial amounts of NO_x (Jain *et al.*, 2014).

3.3. Monthly variation of NO₂

The monthly variation of NO₂ from 2005 to 2022 is depicted in Fig.5 The NO₂ concentration remains relatively flat during the monsoon and post-monsoon seasons. However, a higher peak is observed in the premonsoon season, particularly over Manipur, Mizoram, and Nagaland. During this time, westerly air masses carry NO₂ from the North-west direction, flowing along the Brahmaputra valley and encountering the Arbi-Allong hills in Nagaland [Fig. S1(b)]. Lightning also contributes to enhancing the NO₂ concentration in the region. Additionally, air from Tripura and Bangladesh carries NO₂, leading to higher concentrations over Manipur and Mizoram [Fig. S1(b)]. Tripura exhibits higher NO₂ levels than other states due to local burning and transported NO₂ from Bangladesh [Fig. S1(a)]. In Assam, NO₂ concentration is higher than during the Monsoon and postmonsoon seasons, attributed to local and transported NO₂ from the hilly areas [Fig. S1(a)]. NO₂ is primarily anthropogenic and tends to be highest in highly urbanized areas (Lin et al., 2019). Over the ocean, NO₂ levels are generally low. In China, NO₂ exhibits a seasonal variation, with the highest levels in winter and the lowest in summer. influenced emission by sources and meteorological conditions (Lin et al., 2019). The evolution of aerosol particles in the area is influenced by

both anthropogenic and natural events (Barman *et al.*, 2018). The aerosols in NERI come from various sources, including open agricultural fields, burning vegetation, combustion releases, brick kilns, coal mines, and oil wells (Pathak *et al.*, 2015; Kundu *et al.*, 2018; Barman *et al.*, 2018). The rugged topography in NERI causes aerosols to be mainly confined to the Brahmaputra valley, where air convergence supports an ideal environment for the accumulation of transported and local pollutants (Pathak *et al.*, 2015).

3.4. Natural and anthropogenic sources of TCN

3.4.1. Natural source

This is because higher humidity levels lead to stronger hydrometeor concentration and updraft velocities, both of which contribute to intense lightning. Lightning stands as one of the largest natural sources of atmospheric NOx (von Liebig, 1827; Galloway et al., 2004; Hutchinson, 1954). The high temperature during lightning causes oxygen and nitrogen to combine and form nitric oxide. NO and NO2 are collectively referred to as NOx because NO rapidly reacts with O₃ in the atmosphere, producing NO₂, and achieves equilibrium concerning the photo-dissociation of NO₂ within a few minutes, maintaining the sum of both species essentially unchanged (Bradshaw et al., 1999). LNOx have significant implications for atmospheric chemistry and climate (WMO, 1999; IPCC, 2001). In NERI, lightning activity exhibits bimodal variation, with the first prominent peak (13 flashes/pass) in April and a second lower peak spread between September and October (5.5 flashes/pass), resulting in an average value of 8.4 flashes/pass (Kandalgaonkar, 2010). During the monsoon season, lightning activity is minimal, ranging from 4 to 6 flashes/pass. April and May are considered the most lightning-producing months in the sub-Himalayan region of Nepal (Saha et al., 2012). These observations align with previous studies (Kumar and Kamra, 2012). Seasonal lightning activity over NERI remains one of the major natural sources of NO2. Fig. 6 illustrates the average seasonal variation of lightning flashes across different states in the NERI. The pre-monsoon and monsoon seasons exhibit the highest lightning activity. During the pre-monsoon season, Meghalaya and Tripura record the highest number of lightning flashes, while Arunachal Pradesh experiences the lowest. In the monsoon season, Meghalaya, Tripura, and Mizoram witness the highest lightning frequencies. In the comparison of lightning flashes and NO₂, it is observed that the highest lightning states have not shown the highest concentration of NO₂. It is noteworthy that the NO₂ existence over NER indicates anthropogenic emissions e.g. biomass burning, vehicle emission, and long-range transported NO_2



Fig. 7. Sidewise annual variation of TCN from 2005 to 2022

3.4.2. Anthropogenic source

Anthropogenic NO₂ primarily originates from combustion engines that burn fossil fuels, such as oil, natural gas, and coal, to produce energy. NOx formation during combustion processes involves three gas phase reaction mechanisms (García et al., 2014): thermal NOx mechanism (resulting from atmospheric nitrogen oxidation at high temperatures >1300 °C), fuel NOx mechanism (involving oxidation of fuel-bound nitrogen), and prompt NOx mechanism (caused by the reaction of CHi-radicals with atmospheric nitrogen in the flame front). For fossil fuel combustion, thermal and prompt NOx mechanisms play a significant role. However, in combustion, the combustion chamber's biomass temperature typically remains below 1300 °C, leading to the dominance of the fuel-N mechanism in NOx formation. Biomass burning can occur naturally or be initiated by human activities such as agriculture (crop residue burning), cooking, and heating. People often utilize solid biomass as a renewable and carbon-neutral energy source, replacing fossil fuels. Despite its environmental benefits, there are concerns regarding biomass combustion in small-scale domestic appliances. In the NERI region, the principal sources of NO_2 are attributed to an increasing number of vehicles and factories, biomass mass burning in household appliances, and the burning of crop residues (Fig. 2).

3.4.3. LULC changes and forest fire

India's North Eastern Region (NER) has some of the largest tropical and sub-tropical forest reserves of wet evergreen, semi-evergreen, moist deciduous, coniferous forests, mixed forests, and shrubs (Roy and Joshi, 2002), covering almost 64.66% of its geographic area. Every year forest cover of NERI undergoes significant changes due to shifting (called "Jhum") cultivation. Some people burn forest areas to create new agricultural fields, leading to fires from those fields (Puri et al., 2011) and causing forest fires. Historical data analysis of forest fire counts in NERI shows an average of 100 thousand fire events annually (Sarma et al., 2017). In the FSI, 2021; a decrease of 1020 sq. km of forest area in the region has also been reported due to shifting cultivation, natural calamities, development activities, etc. Spatial variation in peak fire months or seasons was observed over the Indian subcontinent in the study by Chand et al. (2006) which was attributed to the varying forest types and climatic conditions throughout the region.



Fig. 8. Variations in TCN, fire counts and LULC types over the NER

For example, the same study stated that they reported intense fire episodes over Himalayan zone forests during May and June every year due to moist conditions in April and prevailing dry weather conditions in May. Whereas, high forest fires were recorded in the tropical evergreen forests of North East India during February, March, and April with a maximum number of ~1200 fires per day. Kharol et al. (2008) also recorded high forest fire incidents over NER of India due to the burning of agricultural lands in March and April. In the same study, the author also mentions high forest fire incidents in Mizoram due to forest fires. Fire counts vary from 64,092 to 1, 07,506 in March, 4,972 to 6,204 in January, and 12,268 to 28,393 in February. From 2009 to 2011 there was a maximum fire count in winter and the third maximum fire count in pre-monsoon, related to the drought episodes (NRAA, 2009, 2013). It is known that during drought, less availability of moisture in vegetation makes it more flammable, thereby increasing the probability of forest fire (Littell et al., 2016).

The LULC changes over the north-eastern region of India (Fig. 8) showed an increase in forest cover from

2005-2006 to 2010-2011, primarily due to regeneration in abandoned shifting cultivation areas and afforestation activities (ISFI-FSI 2007). This trend was followed by a decline in forest cover towards 2015-2016 (ISFI-FSI 2013), attributed to factors such as biotic pressure, soil erosion, and ongoing shifting cultivation practices (ISFR-FSI 2021). Agricultural land showed a slight decrease during this period, likely due to land degradation resulting from unsustainable agricultural practices. This degradation may have contributed to an increase in barren land, potentially caused by soil erosion or the abandonment of fields (Singh & Chaudhary, 2023).

The increase in forest density during this period, coupled with episodes of drought (NRAA, 2009, 2013), likely contributed to the high fire counts observed in the winter and pre-monsoon seasons of 2009 - 2011 (Borgohain *et al.*, 2023). Reduced moisture availability in vegetation, combined with the accumulation of dry biomass from denser forests, likely provided ample fuel for forest fires, thereby elevating the risk of fire outbreaks in the region (Littel *et al.*, 2016).



Fig. 9. TCN and TCO over NEI states from 2005-2019

Simultaneously, the rise in built-up areas points to increased urbanization and anthropogenic activities, which are associated with elevated NO₂ levels, a pollutant primarily originating from human activities (Lin *et al.*, 2019). The decline in vegetation cover, indicated by the increase in barren land, may have reduced the region's capacity to act as a sink for pollutants like NO2, leading to its accumulation (Gourdji, 2018).

3.4.4. TCN and Tropospheric Ozone

Tropospheric ozone (O₃) is a short-lived climate pollutant formed through the interaction of sunlight with volatile organic compounds (VOCs) and nitrogen oxides (NO_x), which are mainly produced by human activities. O₃ and NO_x, consisting primarily of nitric oxide (NO) and nitrogen dioxide (NO₂), are dynamically interconnected. Nitric oxide reacts with ozone to produce NO₂, establishing a near-equilibrium state within minutes through the photodissociation of NO₂, which stabilizes the combined concentrations of NO and NO₂ (Bradshaw *et al.*, 1999). This equilibrium is why NO and NO₂ are collectively referred to as NO_x. Sources of NO_x include soil emissions, fossil fuel combustion, biomass burning, and lightning, all of which significantly contribute to atmospheric NO₂ levels (Richter *et al.*, 2005; Lin, 2012). The chemical relationship between O₃ and NO_x is central to atmospheric chemistry, as their concentrations are interrelated (Clapp & Jenkins, 2001). This interplay underscores the role of NO₂ as an indicator of ozone levels and the complexity of tropospheric composition.

The correlation between monthly averaged TCN and troposphere column O_3 (TCO) over the different seasons in the 2005-2019 periods varies across north-eastern Indian states due to differences in emission sources, meteorological conditions, and topography (Fig. 9). In general, winter (Win) and pre-monsoon (PrM) seasons exhibit positive correlations, with the strongest seen in Manipur (r = 0.41) and Nagaland (r = 0.57) during these periods, indicating that higher NO₂ corresponds with increased ozone. Negative or near-zero correlations are more common during monsoon (Mon) and post-monsoon (PsM), as seen in Meghalaya (r = -0.31, PsM) and Mizoram (r = -0.16, PsM), likely due to reduced ozone



Fig. 10. Frequency analysis of NO₂ Concentrations in NERI

formation from wet deposition and cloud cover. Sikkim and Arunachal in the mountainous terrains show consistently weak correlations across all seasons, while others like Assam and Mizoram have mixed results, with both positive and negative associations depending on the season.

3.5. Frequency distribution and air quality

The frequency distribution of the number of days with different levels of TCN concentrations throughout the year is depicted in the figure for all states in the NERI (Fig 10). TCN refers to the amount of nitrogen present in the entire atmospheric column above a specific location, measured in units of mol cm⁻². This parameter is crucial for understanding air quality and pollution levels, as nitrogen compounds play a significant role in various atmospheric processes and can have both natural and anthropogenic sources. The figure shows that Tripura and Assam consistently experience more than 50 percent of

TCN values days each year with exceeding 1×10^{15} mol cm⁻². This indicates that these states often have high concentrations of nitrogen compounds in the atmosphere, which can be attributed to various sources, such as industrial activities, vehicular emissions, and agricultural practices. On the other hand, Meghalaya, Mizoram, and Nagaland have more than 25 percent of days with TCN values ranging between 0.75×10^{15} and $1{\times}10^{15}$ mol cm^-2. Manipur, on the other hand, has approximately 10 percent of days with a similar TCN concentration range. These states exhibit moderately elevated TCN levels, suggesting a moderate impact on air quality, likely due to a combination of both local emissions and atmospheric transport of pollutants from neighboring regions. The findings from this frequency distribution analysis provide valuable insights into the air quality status of the NERI states. The higher TCN concentrations in Tripura and Assam imply that these areas might require specific attention and intervention to address air pollution concerns Similarly, the moderate.



Fig. 11. Surface NO2 and TCN over CPCB stations

TCN concentrations observed in Meghalaya, Mizoram, Nagaland and Manipur call for continued monitoring and mitigation efforts to maintain and improve air quality in these regions. Overall, this scientific analysis highlights the importance of understanding and managing nitrogenrelated air pollution to safeguard the environmental health and well-being of the NERI states.

Surface NO₂ over the years 2019-2023 were collected from the 18 CPCB monitoring stations over the 8 NE states and its average is compared with average TCN data over the same locations and over the same time period. This data is represented in Fig. 11, showing the variation of these concentrations over the different stations. In Assam, the oil refinery, paper, and cement industries are major contributors to air pollution (Pandey & Ghosh, 2000), showing relatively higher emissions of ground level NO₂. Byrnihat exhibits significantly higher surface NO₂ levels compared to other stations, despite having TCN values similar to nearby locations in Assam, potentially indicating substantial local emissions with subsequent dispersion into surrounding areas. The Byrnihat Industrial Area, located near the Assam-Meghalaya border, has been recognized as a critically polluted region by the CPCB over recent years (Borgohain et al., 2024; CPCB, 2020). Conversely, stations in Tripura report the highest TCN values among all monitored locations but show moderate surface NO₂ levels, likely due to emissions from non-ground sources or the transport of pollutants from nearby regions (Borgohain et al., 2023).



Fig. 12. NO₂ concentration over different states of NERI during and pre and post-lockdown pre-monsoon seasons

3.6. NO₂ rise after COVID-19

In Fig. 12 NO₂ concentration is shown over the various states of NERI during and pre and post-lockdown years of pre-monsoon season. In pre-monsoon-20, AR, AS, ML, MZ, TR, and SK show a lower NO₂ concentration compared to pre-monsoon-19 (Table1). At the same time, MN and NG showed a higher NO₂ during pre-monsoon-20 compared to pre-monsoon-19.

nest	Regio
and	analys

820

NO2 concentration over different states of NERI during and pre and post-lockdown pre-monsoon seasons

TABLE 1

State —	Pre-monsoon-19	Pre-monsoon-20	Pre-monsoon-21
		NO ₂ (mol cm ⁻²)	
AR	8.30×10 ¹⁴	8.04×10^{14}	9.33×10 ¹⁴
AS	1.98×10^{14}	1.94×10 ¹⁵	2.39×1015
MN	2.75×10 ¹⁵	2.79×10 ¹⁵	3.35×1015
MG	2.01×10 ¹⁵	1.88×10 ¹⁵	2.36×1015
MZ	2.58×10 ¹⁵	2.44×10 ¹⁵	2.99×1015
NG	1.88×10 ¹⁵	1.93×10 ¹⁵	2.43×1015
SK	4.77×10 ¹⁵	4.45×10 ¹⁴	5.86×10 ¹⁴
TR	2.28×10 ¹⁵	2.04×10 ¹⁵	2.40×10 ¹⁵

In pre-monsoon-21, NO₂ concentration increased in all states. In contrast, the rise of NO2 in AS, ML, MN, NG, and TR are significant. A decrease in NO₂ concentrations has been reported as a result of lockdown measures to reduce the spread of COVID-19 (Koukouli et al., 2021; Field et al., 2020; Bauwens et al., 2020; Liu et al., 2020; Prunet et al., 2020). Questions remain, however, regarding the relationship of satellite-derived atmospheric column NO₂ data with health-relevant ambient ground-level concentrations, and the representativeness of limited ground-based monitoring data for global assessment. Cooper et al. (2022) reported with the help of TROPOMI satellite data (during 2019-2020) that NO₂ changes in more than 200 cities, including 65 cities without available ground monitoring, largely in lower-income regions. Mean country - level population - weighted NO2 concentrations are $29\% \pm 3\%$ lower in countries with strict lockdown conditions than in those without (Cooper et al., 2022). Relative to long-term trends, NO₂ decreases during COVID-19 lockdowns exceed recent Ozone Monitoring Instrument (OMI)-derived year-to-year decreases from emission controls, comparable to 15 ± 4 years of reductions globally (Cooper et al., 2022).

4. Conclusions

In summary, this comprehensive 17-year analysis of TCN concentrations in the NERI has revealed seasonal and long-term trends, shedding light on the dynamics of air quality in this region. TCN levels exhibited distinct variations across the four seasons, with pre-monsoon and winter months consistently displaying the highest concentrations, primarily due to biomass burning and increased anthropogenic activities. The long-term trends in TCN demonstrated an overall increase, reflecting the growing impact of factors like rising vehicle numbers, industrial expansion and population density.

Monthly variation of NO₂ concentrations highlights the significance of the pre-monsoon season, where various factors, including lightning and transported NO₂, contribute to elevated NO₂ levels. Both natural and anthropogenic sources, such as forest fires, biomass burning, and combustion engines, have been identified as key contributors to TCN concentrations. LULC changes in relation to TCN concentrations across different states highlight the effect land cover effects on air quality. Troposphere ozone, as a climate pollutant, was compared with TCN data to explore the interactions between atmospheric pollutants. Frequency distribution analysis of air quality status of NERI states emphasized the need for targeted interventions in areas where elevated TCN levels are observed.

The study also explored the impact of the COVID-19 pandemic, revealing fluctuations in NO₂ concentrations, with some states experiencing lower levels during lockdowns in pre-monsoon-20, followed by increased concentrations in pre-monsoon-21.

This study highlighted the various factors influencing the air quality in NERI and calls for informed measures to address evolving air quality challenges in the face of increasing anthropogenic influences and natural variations. The study offers valuable insights into tropospheric column nitrogen dioxide in the North Eastern Region of India (NERI), with a comprehensive 17-year analysis highlighting seasonal and long-term trends. The study effectively integrates data on both natural and anthropogenic sources and examines the impact of the COVID-19 pandemic on NO_2 levels. Its focus on the region's specific environmental conditions and the combination of satellite and ground-based data strengthen the findings.

Future research should focus on several key areas, including expanding ground-based monitoring to improve the validation of satellite data, assessing the health impacts associated with variability in NO_2 levels, conducting detailed source apportionment studies, exploring how climatic variables affect TCN and NO_2 , and evaluating the impact of air quality regulations. Addressing these areas will enhance our understanding and inform more effective environmental management strategies.

Authors' Contributions

Arup Borgohain: Conceptualization, methodology, manuscript writing, data curation, supervision.

Arban S. Youroi: Data collection and analysis, formal analysis, manuscript writing, visualization.

Rohit Gautam: Writing manuscript and editing, formal analysis.

Manasi Gogoi: Data collection, formal analysis, visualization.

Ribanda Marbaniang: Data collection, formal analysis, visualization

Nilamoni Barman: Data analysis, visualization.

Arundhati Kundu: Data collection, visualization.

Abhay Srivastava: Data collection, visualization.

Shyam S. Kundu: Review and supervision.

S.P. Aggarwal: Review and supervision.

Disclaimer: The contents and views presented in this research article/paper are the views of the authors and do not necessarily reflect the views of the organizations they belongs to.

Reference

- Achakulwisut, P., Brauer, M., Hystad, P. and Anenberg, S. C., 2019, "Global, national, and urban burdens of pediatric asthma incidence attributable to ambient NO₂pollution: estimates from global datasets", *Lancet Planet. Health 3*, e166-e178.
- Anenberg, S. C., Henze, D. K., Tinney, V., Kinney, P. L., Raich, W., Fann, N., ... & Kuylenstierna, J. C., 2018, "Estimates of the global burden of ambient PM 2.5, ozone, and NO 2 on asthma incidence and emergency room visits" *Environmental health perspectives*, **126**, 10, 107004.
- Badarinath, K. V. S., Kharol, S. K., Chand, T. K., Parvathi, Y. G., Anasuya, T. and Jyothsna, A. N., 2007, "Variations in black carbon aerosol, carbon monoxide and ozone over an urban area of Hyderabad, India, during the forest fire season", *Atmospheric Research*, 85, 1, 18-26.

- Barman, Nilamoni, Borgohain, Arup, Raju, P. & Kundu, Shyam Sundar, 2018, "Investigation of Curvature Effect of Ångström Exponent to Classify the Aerosol Types over the Region of Interest (88° 98 °E and 20° -30 °N), Atmospheric Pollution Research, 10. 10.1016/j.apr.2018.09.002.
- Barnes, E. A. and Fiore, A. M., 2013, "Surface ozone variability and the jet position: Implications for projecting future air quality", *Geophysical Research Letters*, **40**, 11, 2839-2844.
- Bauwens, M., Compernolle, S., Stavrakou, T., Müller, J. F., Van Gent, J., Eskes, H., ... & Zehner, C., 2020, "Impact of coronavirus outbreak on NO₂ pollution assessed using TROPOMI and OMI observations. *Geophysical Research Letters*, 47, 11, e2020GL087978.
- Biswal, A., Singh, V., Singh, S., Kesarkar, A. P., Ravindra, K., Sokhi, R. S., ... & Mor, S., 2021, "COVID-19 lockdown-induced changes in NO 2 levels across India observed by multi-satellite and surface observations", *Atmospheric Chemistry and Physics*, 21, 6, 5235-5251.
- Borgohain, A., Gogoi, M., Barman, N., Kundu, A., Banik, T., Kundu, S. S., ... & Aggarwal, S. P., 2023, "Impact of Biomass Burning on Black Carbon and NO₂ Over North Eastern Region of India Using Multi-satellite Observations", *Journal of the Indian* Society of Remote Sensing, 1-13.
- Borgohain, D., Lanong, S.and Jaishi, H. P., 2024, "Heavy metal contamination and health risks in ground water at Byrnihat industrial area: urgent need for remediation and public health safeguards", *Proceedings of the Indian National Science Academy*, 1-12.
- Bradshaw, J., Davis, D., Crawford, J., et al., 1999, "Photofragmentation two-photon laser-induced fluorescence detection of NO₂ and NO: Comparison of measurements with model results based on airborne observations during PEM-Tropics A", *Geophys. Res. Lett.*, 26, 471-474.
- Brook, J. R., Burnett, R. T., Dann, T. F., Cakmak, S., Goldberg, M. S., Fan, X., & Wheeler, A. J., 2007, "Further interpretation of the acute effect of nitrogen dioxide observed in Canadian timeseries studies", *Journal of exposure science & environmental epidemiology*, **17**, 2, S36-S44.
- Borgohain, A., Gogoi, M., Barman, N., Kundu, A., Banik, T., Kundu, S. S., ... & Aggarwal, S. P., 2023, "Impact of Biomass Burning on Black Carbon and NO₂ Over North Eastern Region of India Using Multi-satellite Observations", *Journal of the Indian* Society of Remote Sensing, 1-13.
- Borgohain, D., Lanong, S.and Jaishi, H. P., 2024, "Heavy metal contamination and health risks in ground water at Byrnihat industrial area: urgent need for remediation and public health safeguards", *Proceedings of the Indian National Science Academy*, 1-12.
- Bradshaw, J., Davis, D., Crawford, J., Chen, G., Shetter, R., Müller, M., ... & Sandholm, S., 1999, "Photofragmentation two-photon laser-induced fluorescence detection of NO2 and NO: Comparison of measurements with model results based on airborne observations during PEM-Tropics", A. *Geophysical Research Letters*, 26, 4, 471-474.
- Brook, J. R., Burnett, R. T., Dann, T. F., Cakmak, S., Goldberg, M. S., Fan, X., & Wheeler, A. J., 2007, "Further interpretation of the acute effect of nitrogen dioxide observed in Canadian timeseries studies", *Journal of exposure science & environmental epidemiology*, **17**, 2, S36-S44.
- Bucsela, E. J., Celarier, E. A., Wenig, M. O., Gleason, J. F., Veefkind, J. P., Boersma, K. F.andBrinksma, E. J., 2006, "Algorithm for

NO/sub 2/vertical column retrieval from the ozone monitoring instrument", *IEEE Transactions on Geoscience and remote sensing*, **44**, 5, 1245-1258.

- Celarier, E. A., Brinksma, E. J., Gleason, J. F., Veefkind, J. P., Cede, A., Herman, J. R., ... & Levelt, P. F., 2008, "Validation of Ozone Monitoring Instrument nitrogen dioxide columns", *Journal of Geophysical Research: Atmospheres*, **113**(D15).
- Chand, T. K., Badarinath, K. V. S., Prasad, V. K., Murthy, M. S. R., Elvidge, C. D. and Tuttle, B. T., 2006, "Monitoring forest fires over the Indian region using Defense Meteorological Satellite Program-Operational Linescan System nighttime satellite data", *Remote Sensing of Environment*, **103**, 2, 165-178.
- Cheng, M.M. & Jiang, Hong & Guo, Zheng, 2012, "Evaluation of longterm tropospheric NO₂ columns and the effect of different ecosystem in Yangtze River Delta", *Procedia Environmental Sciences*,13, 1045-1056. 10.1016/j.proenv.2012.01.098.
- Clapp, L. J. and Jenkin, M. E., 2001, "Analysis of the relationship between ambient levels of O₃, NO₂ and NO as a function of NOx in the UK", *Atmospheric Environment*, **35**, 36, 6391-6405.
- Cooper, Matthew & Martin, Randall & Hammer, Melanie & Levelt, P. & Veefkind, Pepijn & Lamsal, L. & Krotkov, Nickolay & Brook, Jeffrey & Mclinden, Chris, 2022, "Global fine-scale changes in ambient NO₂ during COVID-19 lockdowns", *Nature*, **601**, 380-387. 10.1038/s41586-021-04229-0.
- CPCB Central Pollution Control Board. Comprehensive Environmental Pollution Abatement Action Plan Byrnihat Industrial Cluster-Assam. Pollution Control Board, Assam 2020 (2020).
- Crouse, D. L., Peters, P. A., Villeneuve, P. J., Proux, M. O., Shin, H. H., Goldberg, M. S., ... & Burnett, R. T. (2015). Within-and between-city contrasts in nitrogen dioxide and mortality in 10 Canadian cities; a subset of the Canadian Census Health and Environment Cohort (CanCHEC). *Journal of exposure science* & environmental epidemiology, 25, 5, 482-489.
- Field, R. D., Hickman, J. E., Geogdzhayev, I. V., Tsigaridis, K. and Bauer, S. E., 2020, "Changes in satellite retrievals of atmospheric composition over eastern China during the 2020 COVID-19 lockdowns", Preprint at https://doi.org/10.5194/acp-2020-567.
- FSI (Forest Survey of India), 2007, India State of Forest Report 2021. Ministry of Environment, Forest and Climate Change, Government of India.
- FSI (Forest Survey of India), 2013, India State of Forest Report 2021. Ministry of Environment, Forest and Climate Change, Government of India.
- FSI (Forest Survey of India), 2021, India State of Forest Report 2021. Ministry of Environment, Forest and Climate Change, Government of India.
- Fu, F., Purvis-Roberts, K. L. and Williams, B., 2020, "Impact of the COVID-19 pandemic lockdown on air pollution in 20 major cities around the world", *Atmosphere*, 11, 1189.
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., ... & Vöosmarty, C. J., 2004, Nitrogen cycles: past, present, and future. *Biogeochemistry*, 70, 153-226.
- García, R., Pizarro, C., Lavín, A. G.and Bueno, J. L., 2014, "Spanish biofuels heating value estimation. Part II: proximate analysis data", *Fuel*, **117**, 1139-1147.
- Ghude, S. D., Van der, A. R. J., Beig, G., Fadnavis, S. and Polade, S. D., 2009, "Satellite derived trends in NO₂ over the major global

hotspot regions during the past decade and their intercomparison", *Environmental Pollution*, **157**, 6, 1873-1878.

- Gilbert, R. O., 1987, "Statistical Methods for Environmental Pollution Monitoring", Van Nostrand Reinhold, New York.
- Goldberg, D. L., Anenberg, S. C., Griffin, D., McLinden, C. A., Lu, Z., & Streets, D. G. (2020). Disentangling the impact of the COVID-19 lockdowns on urban NO2 from natural variability. *Geophysical Research Letters*, 47, 17, e2020GL089269.
- Gourdji, S., 2018, "Review of plants to mitigate particulate matter, ozone as well as nitrogen dioxide air pollutants and applicable recommendations for green roofs in Montreal, Quebec", *Environmental pollution*, 241, 378-387.
- Hamra, G. B., Laden, F., Cohen, A. J., Raaschou-Nielsen, O., Brauer, M., & Loomis, D., 2015, "Lung cancer and exposure to nitrogen dioxide and traffic: a systematic review and metaanalysis", *Environmental health perspectives*, **123**, 11, 1107-1112.
- Haridas, M. K. M., Gharai, B., Jose, B. and Prajesh, T., 2019, "Multi-year satellite observations of tropospheric NO₂ concentrations over the Indian region", *J. Earth Syst. Sci.*, **128**, 13. https://doi.org/10.1007/s12040-018-1032-2.
- Hutchinson, G. E., 1954, "The biochemistry of the terrestrial atmosphere, in: The Earth as a Planet", edited by: Kuiper, G. P., Univ. Chicago Press, Chicago.
- IPCC: Climate change, 2001, The Scientific Basis, 881 pp., Cambridge University Press, Cambridge, UK, New York.
- Jacob, D. J., 1999, Introduction to atmospheric chemistry. Princeton university press.
- Jain, N., Bhatia, A. and Pathak, H., 2014, "Emission of air pollutants from crop residue burning in India", Aerosol and Air Quality Research, 14, 1, 422-430.
- K. Ravindra, S. Mor, A. Ameena, J. S. Kamyotra and Kaushik, C. P., 2003, "Variation in spatial pattern of criteria air pollutants before and during initial rain of monsoon", *Environmental Monitoring and Assessment*, 87, 2, 145-153.
- Kandalgaonkar, S.S. & Tinmaker, Iqbal & Kanchibhotla, Gayatri, 2010, "A lightning climatology of northeast India:1998-2007", International Journal of Meteorology, 35, 89-98.
- Kharol, S. K., Badarinath, K. V. S. and Roy, P. S., 2008, "Studies on emissions from forest fires using multi-satellite datasets over north east region of India", The international archives of the society for photogrammetry and remote sensing, ISPRS Archives, 37.
- Koukouli, Mariliza, Skoulidou, Ioanna, Karavias, Andreas, Parcharidis, Is, Balis, Dimitrios, Manders, A., Segers, A.J., Eskes, Henk and van Geffen, Jos., 2021, "Sudden changes in nitrogen dioxide emissions over Greece due to lockdown after the outbreak of COVID-19", Atmospheric Chemistry and Physics, 21, 1759-1774. 10.5194/acp-21-1759-2021.
- Kumar, P. R. and Kamra, A. K., 2012, "The spatiotemporal variability of lightning activity in the Himalayan foothills", J. Geophys. Res. Atmos., 117, D24201.
- Kundu, S. S., Borgohain, A., Barman, N., Devi, M. and Raju, P. L. N., 2018, "Spatial variability and radiative impact of aerosol along the Brahmaputra River Valley in India: Results from a campaign", *Journal of Environmental Protection*, 9, 4, 405-430.
- Lin, Chin-An & Chen, Yi-Chun & Liu, Chian-Yi & Chen, Wei-Ting & Seinfeld, John & Chou, C., 2019, "Satellite-Derived Correlation

of SO₂, NO₂ and Aerosol Optical Depth with Meteorological Conditions over East Asia from 2005 to 2015", *Remote Sensing*, **11**, 1738. 10.3390/rs11151738.

- Lin, J.T., 2012, "Satellite constraint for emissions of nitrogen oxides from anthropogenic, lightning and soil sources over East China on a high-resolution grid", *Atmos. Chem. Phys.*, **12**, 2881e2898. http://dx.doi.org/10.5194/acp-12-2881-2012.
- Littell, J. S., Peterson, D. L., Riley, K. L., Liu, Y. and Luce, C. H., 2016, "A review of the relationships between drought and forest fire in the United States", *Global Change Biology*, 22, 7, 2353-2369. https://doi.org/10.1111/gcb.13275.
- Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., ... & Joiner, J. (2020). Abrupt decline in tropospheric nitrogen dioxide over China after the outbreak of COVID-19. *Science Advances*, 6, 28, eabc2992.
- Liu, Q., Malarvizhi, A. S., Liu, W., Xu, H., Harris, J. T., Yang, J., ... & Yang, C. (2021). Spatiotemporal changes in global nitrogen dioxide emission due to COVID-19 mitigation policies. *Science* of the Total Environment, 776, 146027.
- Molina, M. J. and Molina, L. T., 2004, "Megacities and atmospheric pollution", Journal of the Air & Waste Management Association, 54, 6, 644-680.
- Nishanth, T. and Kumar, M. K., 2011, "Increasing trends of tropospheric ozone and NO₂ at the prominent hot spots along the coastal belt of the Arabian Sea in Indian Subcontinent", *International Journal of Environmental Sciences*, 1, 5, 860-870.
- NRAA, 2009, "Drought management strategies-2009", In Draft paper of National Rainfed Area Authority. New Delhi: Ministry of Agriculture, Government of India.
- NRAA, 2013, "Contingency and compensatory agriculture plans for droughts and floods in India-2012", Position paper No. 6 (p. 87). National Rainfed Area Authority, NASC Complex, DPS Marg, New Delhi-110012, India.
- Pandey, R. and Ghosh, S., 2000, "Estimating industrial pollution in India: implications for an effluent charge", Mimeo, National Institute of Public Finance and Policy, New Delhi.
- Pannullo, F., Lee, D., Neal, L., Dalvi, M., Agnew, P., O'Connor, F. M., ... & Sarran, C., 2017, "Quantifying the impact of current and future concentrations of air pollutants on respiratory disease risk in England", *Environmental Health*, 16, 1-14.
- Pathak, Binita, Subba, Tamanna, Dahutia, Papori, Bhuyan, Pradip, Moorthy, K. Krishna, Gogoi, Mukunda, Babu, S., Chutia, Lakhima, Parottil, Ajay, Biswas, Jhuma, Bharali, Chandrakala, Borgohain, Arup, Dhar, Pranab, Guha, Anirban, De, Barin, Banik, Trisanu, Chakraborty, Monti, Kundu, Shyam Sundar, Sudhakar, Sunkari and Singh, Sb., 2015, "Aerosol characteristics in North-East India using ARFINET spectral optical depth measurements", Atmospheric Environment, 125. 10.1016/j.atmosenv.2015.07.038.
- Pathak, Binita and Bhuyan, Pradip, 2015, "Climatology of columnar aerosol properties at a continental location in the upper Brahmaputra basin of North East India: diurnal asymmetry and association with meteorology", Advances in Space Research, 56. 10.1016/j.asr.2015.07.004.
- Prasad, A. K., Singh, R. P. and Kafatos, M., 2012, "Influence of coalbased thermal power plants on the spatial-temporal variability of tropospheric NO₂ column over India", *Environmental Monitoring and Assessment*, **184**, 1891-1907.

- Price, C. and Federmesser, B., 2006, "Lightning-rainfall relationships in Mediterranean winter thunderstorms", *Geophysical Research Letters*, 33, 7.
- Prunet, P., Lezeaux, O., Camy-Peyret, C. and Thevenon, H., 2020, "Analysis of the NO₂ tropospheric product from S5P TROPOMI for monitoring pollution at city scale", *City and Environment Interactions*, 8, 100051.
- Puri, K., Areendran, G., Raj, K., Mazumdar, S.and Joshi, P. K., 2011, "Forest fire risk assessment in parts of Northeast India using geospatial tools", *Journal of forestry research*, 22, 641-647.
- Qie, X., Wei, L., Zhu, K., Qie, K., Xu, C., Sun, Z., ... & Yuan, S., 2022, "Regional differences of convection structure of thunderclouds over the Tibetan Plateau", *Atmospheric Research*, 278, 106338.
- Ranjan, R.and Singh, S., 2017, "Energy deprivation of Indian households: evidence from NSSO data".
- Richter, A., & Burrows, J. P., 2002, "Tropospheric NO2 from GOME measurements", Advances in Space Research, 29, 11, 1673-1683.
- Ropkins, K. and Tate, J. E., 2021, "Early observations on the impact of the COVID-19 lockdown on air quality trends across the UK", *Sci. Total Environ.*,**754**, 142374.
- Roy, P. S. and Joshi, P. K., 2002, "Forest cover assessment in north-east India--the potential of temporal wide swath satellite sensor data (IRS-1C WiFS)", *International Journal of Remote Sensing*, 23, 22, 4881-4896.
- Saha, K., Damase, N. P., Banik, T., Paul, B., Sharma, S., De, B. K. and Guha, A., 2019, "Satellite-based observation of lightning climatology over Nepal", *Journal of Earth System Science*, 128, 1-15.
- Sarma, K. K., Chakraborty, K., Chabukdhara, M., Mondal, P., Borah, S., Sudhakar, S. and Raju, P. L. N., 2017, "Status of forest fire-A case study from northeast India", In Asian conference on remote sensing.
- Schoeberl, M. R., Douglass, A. R., Hilsenrath, E., Bhartia, P. K., Beer, R., Waters, J. W., ... & DeCola, P., 2006, "Overview of the EOS Aura mission", *IEEE Transactions on Geoscience and Remote Sensing*, 44, 5, 1066-1074.
- Sen, P. K., 1968, "Estimates of the regression coefficient based on Kendall's tau", Journal of the American statistical association, 63, 324, 1379-1389.
- Shi, X. and Brasseur, G. P., 2020, "The response in air quality to the reduction of Chinese economic activities during the COVID-19 outbreak", *Geophysical Research Letters*, 47, 11, e2020GL088070.
- Shi, Z., Song, C., Liu, B., Lu, G., Xu, J., Van Vu, T., ... & Harrison, R. M., 2021, "Abrupt but smaller than expected changes in surface air quality attributable to COVID-19 lockdowns", *Science* advances, 7, 3, eabd6696.
- Shi, Z., Tan, Y. B., Liu, Y., Liu, J., Lin, X., Wang, M. and Luan, J., 2018, "Effects of relative humidity on electrification and lightning discharges in thunderstorms", *Terr. Atmos. Ocean. Sci.*, 29, 695-708. https://doi.org/10.3319/TAO.2018.09.06.01.
- Siingh, D., Singh, R. P., Singh, A. K., Kulkarni, M. N., Gautam, A. S. and Singh, A. K., 2011, "Solar activity, lightning and climate", *Surveys in geophysics*, 32, 659-703.
- Singh, V. and Chaudhary, N., 2023, "Land degradation, desertification, and food security in North-East India: Present and future

scenarios", In Sustainable development goals in Northeast India: Challenges and achievements (153-166). Singapore: Springer Nature Singapore.

- Sirois, A., 1998, "A brief and biased overview of time series analysis or how to find that evasive trend", WMO report, (133), 14-18.
- Streets, D. G., Canty, T., Carmichael, G. R., de Foy, B., Dickerson, R. R., Duncan, B. N., ... & Wecht, K. J., 2013, "Emissions estimation from satellite retrievals: A review of current capability", *Atmospheric Environment*, **77**, 1011-1042.
- Tao, Y., Mi, S., Zhou, S., Wang, S. and Xie, X., 2014, "Air pollution and hospital admissions for respiratory diseases in Lanzhou, China", *Environ. Pollut.*, 185, 196-201.
- Tariq, S., Ali, M., Mahmood, K., Batool, S. A. and Rana, A. D., 2014, "A study of tropospheric NO₂ variability over Pakistan using OMI data", *Atmospheric Pollution Research*, 5, 4, 709-720.
- Theil, H., 1950, "A rank invariant method of linear and polynomial regression analysis, part 3", Netherlands Akademie van Wettenschappen. No.53:1397-1412.
- Ul-Haq, Z., Tariq, S., & Ali, M. (2015). Tropospheric NO2 trends over South Asia during the last decade (2004–2014) using OMI data. Advances in Meteorology, 2015(1), 959284.
- Uno, I., He, Y., Ohara, T., Yamaji, K., Kurokawa, J. I., Katayama, M., ... & Burrows, J. P., 2007, "Systematic analysis of interannual and seasonal variations of model-simulated tropospheric NO₂ in Asia and comparison with GOME-satellite data", *Atmospheric Chemistry and physics*, 7, 6, 1671-1681.
- Venter, Z. S., Aunan, K., Chowdhury, S. and Lelieveld, J., 2020,"COVID-19 lockdowns cause global air pollution declines", *Proc. Natl Acad. Sci.*,**117**, 18984-18990.

- von Liebig, J., 1827, "Une note sur la nitrification", Ann. Chem. Phys., 35, 329-333.
- Wang, W., Chen, Y., Becker, S. and Liu, B., 2015, "Variance correction prewhitening method for trend detection in autocorrelated data", *Journal of Hydrologic Engineering*, 20, 12, 04015033.
- Wang, X. L. and Swail, V. R., 2001, "Changes of extreme wave heights in Northern Hemisphere oceans and related atmospheric circulation regimes", *Journal of Climate*, 14, 2204-2221.https://doi.org/10.1175/1520-0442(2001)014.
- WMO, 1999, Scientific Assessment of Ozone Depletion: 1998, World Meteorological Organisation, Geneva, Switzerland.
- Yoo, Jung-Moon & Lee, Yu-Ri & Kim, Dongchul & Jeong, Myeong-Jae & Stockwell, William & Kundu, Prasun & Oh, Soo-Min & Shin, Dong-Bin & Lee, Suk-Jo., 2014, "New indices for wet scavenging of air pollutants (O₃, CO, NO₂, SO₂ and PM₁₀) by summertime rain", *Atmospheric Environment*, 82, 226-237.
- Yue, S., Pilon, P. and Cavadias, G., 2002, "Power of the Mann-Kendall and Spearman's rho tests for detecting monotonic trends in hydrological series", *Journal of hydrology*, 259, 1-4, 254-271.
- Zeng, W., Zhao, H., Liu, R., Yan, W., Qiu, Y., Yang, F., ... & Zhan, Y., 2020, "Association between NO₂ cumulative exposure and influenza prevalence in mountainous regions: a case study from southwest China", *Environmental research*, **189**, 109926.
- Ziemke, J. R., Chandra, S., Duncan, B. N., Froidevaux, L., Bhartia, P. K., Levelt, P. F. and Waters, J. W., 2006, "Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's Chemical Transport Model", *Journal of Geophysical Research: Atmospheres*, **111**(D19).